

Effect of Post-Harvest Handling Practices, Storage Technologies and Packaging Material on Post-Harvest Quality and Antioxidant Potential of *Solanum Aethiopicum* (Shum) Leafy Vegetable

Sekulya S.¹, Nandutu A.¹, Namutebi A.², Ssozi J.², Masanza M.³, Jagwe J. N.⁴, Kasharu A.⁵, Rees D.⁶, Kizito E. B.³, Acham H.^{2,*}

¹Department of Biochemistry, College of Natural Sciences, Makerere University, Kampala, Uganda

²Department of Food Technology and Nutrition, College of Agricultural and Environmental Sciences, Makerere University, Kampala, Uganda

³Department of Agricultural and Biological Sciences, Faculty of Science and Technology, Uganda Christian University, Mukono, Uganda

⁴Farmgain Africa Limited, Kampala, Uganda

⁵Coalition for Health Agricultural Income Networks, Kampala, Uganda

⁶Natural Resources Institute, University of Greenwich, Chatham Maritime, Kent ME4 4TB, UK

*Corresponding author: hacham@caes.mak.ac.ug

Abstract Several studies have supported the use of vegetables as foods as well as medicinal plants. However, most especially for the leafy types of vegetables, their high moisture content gives them a short shelf life. On average *Solanum aethiopicum* (Shum) has a shelf life of one day, making it unable to keep fresh for a long time. The objective of this study was to determine the effect of post-harvest handling practices and storage technology on the post-harvest quality and antioxidant activity in *S. aethiopicum*, as well as determine the packaging material that could be able to maintain a high post-harvest quality during storage. The post-harvest handling and storage technologies were tested under three experimental conditions. Experiment one involved placing 2.0 kg of the harvested *S. aethiopicum* with roots intact (RI) and others with roots cut-off (RC) in a charcoal cooler (-CC), 21.0±1.00 °C, 95.67±3.01 %rh; in ambient storage (-AC), 23.8±2.86 °C, 69.38±6.72 % rh; and in cold room (-CR), 7.17±1.30 °C, 95.80±3.19 %rh. Experiment two involved storing 2.0 kg of *S. aethiopicum* in charcoal cooler with no water treatment (TT⁻) and in ambient storage while immersing in portable water for 2 to 3 seconds during the day (TT⁺). Experiment three involved packing 1.0 kg of *S. aethiopicum* sample of both RC and RI state to assess the effectiveness of the packaging materials (0.1 cm meshed perforated polyethylene (RC0.1), 0.5 cm meshed perforated polyethylene (RC0.5) and a 60 µm perforated polyethylene (RC60µm) in maintaining quality of the vegetables. The edible parts of the vegetable were tested for moisture content, percentage weight loss, chlorophyll content, polyphenol content and total antioxidant activity (as measures of post-harvest quality and shelf life) after every 24 hours. The antioxidant activity was determined by screening for free radical scavenging properties using diphenyl picryl hydrazyl (DPPH), Ferric reducing antioxidant power (FRAP) and ascorbic acid as standard. The results revealed that Shelf life was found to increase (from one day to four days) when the vegetable was intermittently immersed in portable water for 2 to 3 seconds after every one hour during the day for vegetables in ambient storage both with roots intact (RI(TT⁺)-AC) and with roots cut-off RC(TT⁺)-AC). The samples stored in cold room and charcoal cooler showed slow and comparable reduction (percent) of weight for both intact and roots cut. The chlorophyll content decreased in all storage conditions, with ambient conditions showing the most rapid decrease. The total polyphenol fluctuated within relatively small limits for both with intact and roots cut-off when stored in cold room and charcoal cooler (6.25±0.05 to 9.35±0.05 mgGAE/gfw; respectively) within the four days of storage. Storage in ambient conditions indicated an increase in total polyphenol content from 9.35±0.05 to 14.77±0.12 mgGAE/gfw for that with roots intact (RI-AC) and to 13.65±0.06 mgGAE/gfw for roots cut-off (RC-AC). The increase in total polyphenol content in the ambient storage led to increased total antioxidant activity compared to that stored in cold room and charcoal cooler that remained almost constant. The 60 µm perforated polyethylene and 0.1 cm meshed perforated polyethylene retained more moisture (84.55±0.18 % and 85.20±0.03 %; respectively) and showed minimal percentage of weight loss (9.69±0.25 %) with the highest chlorophyll content (8.06±0.02 mg/g dwb) on day four when stored in the charcoal cooler, making it the best tested packaging material.

Keywords: *Solanum aethiopicum*, charcoal cooler, shelf life, total polyphenol, total antioxidant activity, Ferric reducing antioxidant power (FRAP), diphenylpicrylhydrazyl (DPPH), perforated polyethylene

Cite This Article: Sekulya S., Nandutu A, Namutebi A., Ssozi J., Masanza M., Jagwe J. N., Kasharu A., Rees D., Kizito E. B., and Acham H., “Effect of Post-Harvest Handling Practices, Storage Technologies and Packaging Material on Post-Harvest Quality and Antioxidant Potential of *Solanum Aethiopicum* (Shum) Leafy Vegetable.” *American Journal of Food Science and Technology*, vol. 6, no. 4 (2018): 167-180. doi: 10.12691/ajfst-6-4-6.

1. Introduction

The Solanaceae vegetable family has over 1000 species worldwide with at least 100 indigenous species in Africa [1]. Several studies have supported use of these vegetables as foods and medicinal plants [1], but despite their high nutrient content and benefits these come with, there has been limited research on local vegetable varieties in Uganda [2]. *Solanum aethiopicum* is the most popular of the Solanaceae leafy vegetables grown in east and central Uganda, and consumed in most of the restaurants in Kampala [3]. However, vegetables are sensitive to environmental changes and especially changes to availability of water and temperature [4]. Additionally, post-harvest losses account for 50% of the total vegetable loss in the developing world as a result of inadequate infrastructure, poor technology and over production [5]. Any change in quality due to post-harvest handling in the market therefore leads to losses to the farmer.

Vegetables are also seasonal, with peak production occurring in May, September and October [6]. The highest yields of *S. aethiopicum* Shum group are got under warm humid conditions. The yield is significantly affected by limited water supply during growth, this necessitates growing *S. aethiopicum* during rainy seasons or where there is reliable supply of water [7]. Drought seasons always give low yields as droughts significantly affect the crop physiology [7].

Post-harvest handling practices for *S. aethiopicum* have been documented to be rudimentary [8] which are likely to affect the quality of the vegetables. Some of the several factors that affect post-harvest quality of a vegetable includes; temperature in storage room, percentage relative humidity, light intensity, carbon dioxide concentration [9], cutting which brings about wounding that promotes phytochemical synthesis [10,11], increased light transmission and oxygen concentration which increase chlorophyll loss leading to deterioration [12,13,14]. On average, the shelf life of *S. aethiopicum* has been reported locally to be one day. Uganda is known to be a fruit basket where most of the food is consumed in its fresh form, this calls for development of post-harvest technologies that can prolong the shelf life of harvested leafy vegetables.

One of the most recent innovations has been on drying of *S. aethiopicum* as a processing technology [15]. This had its own limitations in terms of acceptability, where the dried vegetable product was rated lower than the fresh vegetables. Therefore, post-harvest handling must be done in the best way possible in order to increase shelf life and availability of the vegetable in its fresh form. If post-harvest handling is not well done, there is likely to be a loss in terms of the amount and quality of vegetable. Different vegetables require different methods of post-harvest handling that may generally depend on the maturity, water content of the leaves, texture and the shelf life at optimum temperature [16], and the post-harvest

handling method adopted determines the ultimate quality of the vegetable in terms of texture, color, taste, nutrient and phytochemical content [17]. There is need to maintain temperature as low as possible. Currently mechanical refrigerators are used which are energy intensive, unaffordable to local farmers and require constant supplies of electricity, unavailable in rural parts of the country. Despite the losses in quality and quantity, there is insufficient recorded information available regarding the influence of post-harvest handling practices on the moisture content, weight loss, phytochemical content and antioxidant activity of *S. aethiopicum* vegetable in Uganda. The objective of this study was to determine the post-harvest handling practices, storage technology and packaging material that can maintain a high post-harvest quality of *S. aethiopicum* leafy vegetable during storage for longer shelf life. Thus the effect of handling practices, packaging material and storage conditions on physical attributes (moisture content and weight loss) and the phytochemical attributes (total anti-oxidant capacity, total polyphenols and chlorophyll content) of *S. aethiopicum* was determined, by comparing two different storage conditions including use of the charcoal cooler which is affordable by local farmers and the electrical refrigerator.

2. Materials and Methods

2.1. Chemicals

All chemicals and reagents used were of analytical grade, purchased from sigma Germany. The filter papers were purchased from Whatman, UK.

2.2. Procurement of Vegetable Samples

The *S. aethiopicum* vegetables were procured from local farms of Kabbubu, Nangabo Sub County, Wakiso district in central Uganda at maturity and subjected to different post-harvest handling and storage methods. All the analyses were done in the research laboratory of the Food Technology and Nutrition Department, Makerere University, Uganda. The edible parts of the plant (leaves) were removed, washed under running water, crushed and used to extract and determine polyphenols total antioxidant activity and chlorophyll. The samples used for moisture content determination were not washed.

2.3. Experimental Design for Handling Practices and Storage Technologies

Three experimental designs were set up (Figure 1), two for studying handling practices and storage technologies, and the third for suitability of packaging material in extending the shelf life of *S. aethiopicum*. All experiments were done in 3 replicates.

In experiment one, two (2.0) kg of the vegetables were placed in three (3) perforated buckets and then; one was

kept in cold room -CR (7.17 ± 1.30 °C, 95.80 ± 3.19 %rh), one in the fabricated charcoal cooler -CC (21.0 ± 1.00 °C, 95.67 ± 3.01 %rh), and the other kept at ambient conditions -AC (23.8 ± 2.86 °C, 69.38 ± 6.72 %rh). In experiment two, 2.0 kg of the vegetables were placed in two perforated buckets then one was stored in the fabricated charcoal cooler and the other at ambient conditions (23.8 ± 2.86 °C, 69.38 ± 6.72 %rh). For those stored at ambient conditions, their leaves were dipped in portable water (as a treatment - TT+) for 2-3seconds after every one hour for twelve hours each day for five days. In experiment three, one (1.0) kg of the vegetable with roots cut-off (RC) was each packaged in three packaging materials; high gauge ($60 \mu\text{m}$) perforated polyethylene material, 0.5 cm pore size meshed polyethylene material and 0.1 cm pore size meshed (gauze) polyethylene material. One lot was placed in cold room RC-CR ($60 \mu\text{m}$, 0.5, 0.1) (7.17 ± 1.30 °C, 95.80 ± 3.19 %rh);

and the other in the fabricated charcoal cooler RC-CC ($60 \mu\text{m}$, 0.5, 0.1). The vegetables were kept under the respective treatment conditions for five days, with analysis for moisture content, weight loss, chlorophyll content, total polyphenol content and total antioxidant activity carried out each consecutive day.

2.4. Preparation of Extracts for Analyses

Extraction was done following Wissam et al., [18], with slight modifications. The edible parts of fresh leaves were blended and 1.0 g of the paste dissolved in 50 ml of 80 % methanol solution in a conical flask placed in a thermostatic water bath shaker at 45°C for 20 minutes. The liquid extract was separated from solids by centrifugation at 2000rpm for 10minutes and the supernatant stored at -20°C . This extraction was done in 3 replicates.




Figure 1. The study profile

2.5. Determination of Polyphenol Content

The total phenolic content was determined following Folin-Ciocalteu [19]. To 0.5 ml of extract, 2.5 ml of 10 % Folin-Ciocalteu's reagent dissolved in water was added, followed by 2.5 ml of 7.5 % NaHCO₃, incubated at room temperature in the dark for 45 minutes and the absorbance determined at 765 nm. The mean value of the replicate samples for the absorbance was then calculated. The standard solutions of gallic acid of concentrations 0.01, 0.02, 0.03, 0.04, 0.05 mg per ml were used to construct a standard calibration curve as described [20] and the concentration of phenolics were read in mg per g of fresh sample.

2.6. Determination of Antioxidant Activity

2.6.1 Free Radical Scavenging Activity by the DPPH Method

The free radical scavenging activity was assayed following [21], and kept at -20 °C and a 0.1 mM DPPH was prepared by diluting 10 ml of the stock solution with 90 mL of methanol. Ascorbic acid was used as the standard, prepared in concentrations of 25, 50, 75, 100 and 125 µg/ml. Equal volumes of 1.5 ml of the standard and the sample were added, kept in the dark for 30 minutes and absorbance measured at 517 nm using Genesys 10-UV spectrophotometer (Thermo Electron Corporation, Madison WI, USA). The percentage inhibition of both standard and samples was calculated using the formula;

$$\% \text{ inhibition} = \frac{AB - AA}{AB} \times 100$$

Where; AB is absorbance of blank sample and AA is absorbance of sample.

A plot of % inhibition against ascorbic acid concentration was generated to show a calibration curve for the Ascorbic Acid equivalent (mg/g) of fresh sample.

2.6.2 Radical Scavenging Activity by FRAP Method

Ferric reducing antioxidant power assay was performed following Iqbal, Salim, and Lim [22], with slight modifications. To the extract (1 ml) was added 1 ml of FRAP reagent, that was prepared by mixing of 300 mM sodium acetate buffer (pH 3.6), 10 mM 2,4,6-tri (2-pyridyl)-s-triazine (TPTZ) solution and 20 mM FeCl₃.6H₂O in a ratio of 10:1:1, and diluted with water to a total volume of 4 ml. Ascorbic acid, used as the standard was prepared in concentrations of 25, 50, 75, 100 and 125 µg/ml following [23]. The mixture was incubated in a water bath at 37°C for 30 minutes and absorbance determined at 593 nm using Genesys 10-UV spectrophotometer (Thermo Electron Corporation, Madison WI, USA). The results were expressed as ascorbic acid equivalent in mg per g of fresh sample.

2.7. Determination of Chlorophyll

Acetone was used to extract chlorophyll, which was later determined using a UV Vis spectrophotometer [24,25]. Accurately weighted 0.1 g of fresh leaf sample was taken, and macerated in 10 ml of 80 % acetone

solution using celite. The mixture was filtered using whatman No 1 filter paper; and the filtrate diluted with 80 % acetone to make 100 ml. The mixture was then analyzed for Chlorophyll-a and Chlorophyll-b content in a UV-spectrophotometer at 663.2 nm and 646.8 nm using Genesys 10-UV spectrophotometer (Thermo Electron Corporation, Madison WI, USA). The quantification of chlorophyll a and b was done using the equation by [24] and results expressed in mg/g dwb.

$$Cha = 12.25A_{663.2} - 2.79A_{646.8} \quad (1)$$

$$Chb = 21.5A_{646.8} - 5.1A_{663.2} \quad (2)$$

2.8. Determination of Moisture Content

The moisture content was determined as described in AOAC [26]. A thoroughly washed Petri-dish was placed in the oven to dry and then weighed. A sample of 3.0 g of blended *S. aethiopicum* leaves was placed in the weighed Petri dish, and then placed in an oven to dry at 60°C for 16 hours. The dish and dry sample were transferred to a desiccator to cool to room temperature before being weighed again. Every sample was analysed in 3 replicates.

2.9. Weight Loss

The vegetable was weighed (2.0 kg) on the day of harvest and stored under three different treatment conditions (cold room, charcoal cooler and ambient conditions). After every 24 hours the vegetables were weighed for five days and the percentage weight loss determined using the formula;

$$\frac{W_o - W_t}{W_o} \times 100 \quad (3)$$

Where *W_o* is the weight on day zero and *W_t* is the weight within an interval of 24 hours.

2.10. Control Experiment

All the different parameters including vitamin C, chlorophyll content, moisture content, weight loss, total antioxidant capacity and total polyphenol were determined on the day of harvest (day zero) and the results compared with the results of the subsequent days.

2.11. Determination of Temperature and Humidity

Calibrated data loggers were put in the air space of the charcoal cooler, cold room and stored at ambient conditions to record the temperature and humidity changes after every 30 minutes for five days. The average temperature (°C) and relative humidity (%rh) was determined each day.

2.12. Data Analysis

The data analyzed by using a two-way Analysis of Variance. The significant differences were obtained using the Tukey HSD test (*p* ≤ 0.05). All data was analyzed using SPSS version 16.0 for windows (SPSS, Inc., Chicago, IL, USA).

3. Results

The percentage moisture content of RC-CR decreased from $83.91 \pm 0.18 \%$ to $79.69 \pm 0.69 \%$ on day zero to day four; while RC-CC decreased from $83.91 \pm 0.18 \%$ to $78.95 \pm 0.23 \%$ on day zero to day four and RC-AC decreased from $83.91 \pm 0.18 \%$ to $22.01 \pm 0.08 \%$ on day zero to day four. RI-CR remained almost constant that is from $83.88 \pm 0.08 \%$ to $83.20 \pm 0.01 \%$ on day zero to day four, RI-CC and RI-AC decreased from $83.88 \pm 0.08 \%$ to $80.58 \pm 0.19 \%$ and $83.88 \pm 0.08 \%$ to $40.25 \pm 1.06 \%$ on day zero to day four; respectively. The fastest decrease in percentage moisture content was observed in ambient storage. Both RC-CR, RI-CR, RC-CC and RI-CC showed slight decrease in percentage moisture content from day zero to day four. The data showed that there was variation between the moisture content of RC-CR, RC-CC, RC-AC,

RI-CR, RI-CC and RIAC for each day in the five days of storage, which was significant ($P \leq 0.05$). The percentage weight loss of RC-CR increased from $0.00 \pm 0.00 \%$ to $34.42 \pm 1.63 \%$ on day zero to day four and RI-CR increased from $0.00 \pm 0.00 \%$ to $23.24 \pm 2.4 \%$ on day zero to day four. For RC-CC, the percentage weight loss increased from 0.00 ± 0.00 to $36.22 \pm 1.48 \%$ on day zero to day four for and for RI-CC increased from 0.00 ± 0.00 to $24.44 \pm 0.04 \%$ on day zero to day four. The percentage weight loss of RC-AC increased from $0.00 \pm 0.00 \%$ on day zero to $50.30 \pm 0.33 \%$ on day four and for RI-AC it increased to $42.24 \pm 0.43 \%$ on day four. *S. aethiopicum* RC_AC showed the most rapid increase in the percentage weight loss. There was variation between the percentage weight loss RC-CR, RC-CC, RC-AC, RI-CR, RI-CC and RI-AC for each day in the five days of storage which was significant ($P \leq 0.05$).




Figure 2. Percentage moisture content and percentage weight loss in *S. aethiopicum*, stored in cold room, charcoal cooler and at ambient condition. RI is *S. aethiopicum* with roots intact, RC is *S. aethiopicum* with roots cut-off. RI-CR is roots intact in cold room




Figure 3. Changes in the chlorophyll content of *S. aethiopicum* stored in cold room, charcoal cooler and at ambient condition. RI is *S. aethiopicum* with roots intact, RC is *S. aethiopicum* with roots cut-off. RI-CR is roots intact in cold room, RI-CC is roots intact in charcoal cooler, RI-AC is roots intact in ambient storage, RC-CR is roots cut-off in cold room, RC-CC is roots cut-off charcoal cooler and RC-AC is roots cut-off in ambient storage

The chlorophyll content decreased with duration in varying storage conditions for both *S. aethiopicum* with roots intact (RI) and with roots cut-off (RC). The chlorophyll content of RC-AC and RI-AC decreased from 18.86 ± 0.01 mg/g dwb on day zero to 9.87 ± 0.03 mg/gdwb and 18.86 ± 0.01 mg/gdwb to 11.75 ± 0.03 mg/gdwb respectively. In charcoal cooler, the chlorophyll content decreased from 86.86 ± 0.01 mg/g to 10.29 ± 0.001 mg/g dwb for RC-CC and 18.86 ± 0.01 mg/g dwb to 12.98 ± 0.01 mg/g dwb for RI-CC. For RC-CR, the chlorophyll content decreased from 18.86 ± 0.01 mg/g dwb to 12.37 ± 0.03 mg/g dwb on day four and for RI-CR decreased from 18.86 ± 0.01 to 13.51 ± 0.02 mg/g dwb. The most rapid decrease in chlorophyll content was observed in RC-AC as shown in Figure 2 above. The results showed that there was a statistically significant difference in the chlorophyll content of *S. aethiopicum* kept in the three different storage conditions and between RI and RC in each day of storage ($P \leq 0.05$).

The total polyphenols of RC-CR decreased slightly with duration of storage from 9.35 ± 0.05 to 6.38 ± 0.13 mg GAE/g fw and for RI-CR the total polyphenol content remained relatively constant (Figure 3). The total polyphenol of RC-CC and RI-CC was shown to remain relatively constant. RI-AC and RC-AC showed increase in total polyphenols. However, for RI-AC the total polyphenols remained relatively constant from day zero to day one (6.25 ± 0.05 and 6.30 ± 0.15 mg GAE/g fw) but a significant increase was observed from day two to day five. RI-AC showed the highest increase in total polyphenols on day three and five (15.93 ± 0.26 and 14.77 ± 0.12 mgGAE/g fw respectively) compared to that with RC-AC (13.83 ± 0.08 and 13.65 ± 0.06 mgGAE/g fw respectively) (Figure 6). The results showed that there was a statistically significant difference ($P \leq 0.05$) in the total polyphenols of *S. aethiopicum* stored in the different storage conditions in each day of the storage duration.




Figure 4. Variation in the total polyphenol content of *S. aethiopicum* with storage duration stored when stored in cold room, charcoal cooler and at ambient conditions. RI is *S. aethiopicum* with roots intact, RC is *S. aethiopicum* with roots cut-off. RI-CR is root intact in cold room, RI-CC is roots intact in charcoal cooler, RI-AC is roots intact in ambient storage, RC-CR is roots cut-off in cold room, RC-CC is roots cut-off charcoal cooler and RC-AC is roots cut-off in ambient storage




Figure 5. Variation in the total antioxidant activity (TAA) of the stored *S. aethiopicum* in cold room, charcoal cooler and at ambient condition. RI-FRAP and RC-FRAP are results for FRAP method and RI-DPPH and RC-DPPH are results for DPPH method. RI-CR is roots cut-off in cold room, RC-CC is roots cut-off in charcoal cooler and RC-AC is roots cut-off in ambient storage

RC-CR showed a constant TAA with both FRAP and DPPH methods on day zero and day one. A decrease was observed starting on day two to day four; 3.42 ± 0.03 mgAAE/g fw on day two to 2.3 ± 0.06 mgAAE/g fw on day five when using FRAP method and 4.38 ± 0.12 mgAAE/g fw on day one to 3.49 ± 0.02 mgAAE/g fw on day four when using DPPH method. However, the TAA of RC-CC was observed to remain relatively constant. RI-AC showed a decrease in the TAA for the first two days (2.98 ± 0.03 mgAAE/g fw on day zero to 2.48 ± 0.03 mgAAE/g fw on day one using FRAP method and 4.38 ± 0.03 mgAAE/g fw on day zero to 3.76 ± 0.05 mgAAE/g fw on day one when using DPPH method) and from day two to day four the TAA when determined using the two methods. RC-AC showed an increase in the TAA right on day one and the increase continued to day four when using both FRAP and DPPH method. The results showed that there was a statistically significant difference ($P \leq 0.05$) in the TAA between the storage conditions and that determined for the different days from day zero to day four.

The total polyphenol content of RI(TT⁺)-AC increased from 4.45 ± 0.02 mg GAE/g fw on day zero to 6.56 ± 0.05 mgGAE/gfw on day four and that of RI(TT⁻)-CC increased from 4.45 ± 0.02 mgGAE/g fw to 10.96 ± 0.12 mgGAE/g fw. RC(TT⁺)-AC showed increase in the total polyphenols from 4.51 ± 0.12 mgGAE/g fw on day zero to 7.6 ± 0.05 mgGAE/g fw and that of RC(TT⁻)-CC increased from 4.51 ± 0.12 mgGAE/g fw to 11.98 ± 0.1 mgGAE/g fw (Figure 6). The results showed that there was a slight increase in the total polyphenol content of *S. aethiopicum* RC(TT⁺)-AC. *S. aethiopicum* RC(TT⁺)-AC and RC(TT⁻)-CC showed a faster increase in the total polyphenol content with storage compared with RI(TT⁺)-AC and RI(TT⁻)-CC. This data showed that there was a significant difference ($P \leq 0.05$) in the total polyphenol of RC(TT⁺)-AC, RC(TT⁻)-CC, RC(TT⁺)-AC and RC(TT⁻)-CC from day zero to day four. The FRAP method showed that TAA of *S.aethiopicum* RI(TT⁺)-AC increased from 2.5 ± 0.22 mgAAE/g fw on day zero to 2.72 ± 0.03 mgAAE/g fw on day four and for RI(TT⁻)-CC increased from 2.55 ± 0.15 mgAAE/gfw on day zero to 4.69 ± 0.03 mgAAE/gfw on day four as that of RC(TT⁺)-AC increased from 2.45 ± 0.02 mg AAE/g fw on day zero to 3.34 ± 0.02 mgAAE/g fw on day four but for RC(TT⁻)-CC increased from

2.45 ± 0.02 mgAAE/g fw on day zero to 4.93 ± 0.02 mgAAE/g fw on day four (Figure 7). The DPPH method also showed slight increase in the TAA of *S. aethiopicum* TT⁺. That of RI(TT⁺)-AC increased from 4.32 ± 0.02 mgAAE/g fw to 4.68 ± 0.02 mgAAE/g fw and that of RC(TT⁻)-AC increased from 4.23 ± 0.1 mgAAE/g fw to 4.86 ± 0.03 mgAAE/g fw. The TAA of *S. aethiopicum*, RI(TT⁻)-CC increased from 4.23 ± 0.1 mgAAE/g fw on day zero to 5.21 ± 0.02 mgAAE/g fw and that of RC(TT⁻)-CC increased from 4.32 ± 0.02 mgAAE/g fw on day zero to 5.86 ± 0.02 mgAAE/g fw on day four. The total antioxidant activity of *S. aethiopicum*, TT⁺ increased slightly while TT⁻ remained almost constant when determined using both FRAP and DPPH methods as shown in Figure 6. *S. aethiopicum*, RC showed a more rapid increase in the TAA than RI. The results showed that there was a significant difference ($P \leq 0.05$) between TAA of the *S. aethiopicum*, TT⁺ and TT⁻.

The percentage moisture content of *S. aethiopicum*, RI(TT⁺)-AC increased from 78.91 ± 0.01 % on day zero to 88.70 ± 0.03 % on day four and for RC(TT⁺)-AC, the percentage moisture content increased from 77.28 ± 0.63 % on day zero to 88.77 ± 0.12 % on day four. Percentage moisture content of *S. aethiopicum* RI(TT⁻)-CC decreased from 78.91 ± 0.01 % on day zero to 75.79 ± 0.16 % and that of RC(TT⁻)-CC decreased from 77.28 ± 0.63 % on day zero to 75.81 ± 0.05 % on day four as shown in Figure 7. The difference in the percentage moisture content of *S. aethiopicum*, TT⁺ and TT⁻ was significant at $p \leq 0.05$. The chlorophyll content of *S. aethiopicum*, RI(TT⁺)-AC remained almost constant, that is 10.95 ± 0.04 mg/g dwb on day zero to 10.25 ± 0.01 mg/g dwb on day four while that of RC(TT⁺)-AC decreased from 10.94 ± 0.01 mg/g dwb on day zero to 10.25 ± 0.18 mg/g dwb on day four (Figure 8). The results showed a decrease in the chlorophyll content of *S. aethiopicum*, RI(TT⁻)-CC and RC(TT⁻)-CC. For *S. aethiopicum*, RI(TT⁻)-CC the chlorophyll content decreased from 10.95 ± 0.04 mg/g dwb on day zero to 7.65 ± 0.02 mg/g dwb on day four as for RC(TT⁻)-CC as shown in Figure 9. The chlorophyll content first increased from 10.94 ± 0.01 mg/g dwb on day zero to 12.27 ± 0.16 mg/gdwb on day one then decreased to 6.00 ± 0.03 mg/g dwb on day four. There was a significant difference in the chlorophyll content of *S. aethiopicum* TT⁺ and TT⁻ at $P \leq 0.05$.




Figure 6. Variation in the total polyphenol content of *S. aethiopicum* stored at ambient condition and intermittently immersed in portable water for 2-3 seconds every after one hour during day (TT⁺). RI(TT⁺)-AC is with roots intact in ambient storage, RI(TT⁻)-CC is with roots intact in charcoal cooler, RC(TT⁺)-AC is roots cut-off in ambient and RC(TT⁻)-CC is roots cut-off in charcoal cooler




Figure 7. Effect of immersing *S. aethiopicum* in portable water intermittently for 2-3 seconds after every one hour (TT^+) on antioxidant activity (TAA), under different storage conditions. RI-DPPH and RC-DPPH is TAA determined using DPPH method of *S. aethiopicum* with roots intact and roots cut-off respectively. RI(TT^+)-AC and RC(TT^+)-AC is *S. aethiopicum* in ambient storage with water treatment but with roots intact and roots cut-off respectively. RI(TT^+)-CC and RC(TT^+)-CC is *S. aethiopicum* in charcoal cooler without water treatment but with roots intact and roots cut-off respectively




Figure 8. Effect on percentage moisture content, of immersing harvested *S. aethiopicum* in portable water intermittently but stored at ambient condition (TT^+) and that stored in charcoal cooler but without water treatment (TT^-). RI is with roots intact and RC is with roots cut-off. RI(TT^+)-AC and RC(TT^+)-AC is that with roots intact and roots cut-off respectively in ambient storage with water treatment. RI(TT^-)-CC and RC(TT^-)-CC is that with roots intact and roots cut-off respectively in charcoal cooler without water treatment

The percentage moisture content of *S. aethiopicum* RC-CR60 μm and RC-CC60 μm decreased from 86.91 ± 1.76 % on day of harvest to 85.66 ± 0.06 % and 84.55 ± 0.18 % respectively on day four. The RC-CC0.5 showed the lowest moisture content on day four of 80.55 ± 0.07 % while RC-CR0.1 showed the lowest moisture content on day four of 82.99 ± 0.03 % as shown in Table 1. The results

showed that all the packaging material had high moisture retention within the shoots of *S. aethiopicum*. The RC-CR60 μm polyethylene retained most of the water. The analysis showed that there was a significant difference in the moisture content of the packaged *S. aethiopicum* stored in cold room and charcoal cooler for each day among the three different packaging materials.

Table 1. Moisture content of packaged *S. aethiopicum* stored in cold room and charcoal cooler

Duration (days)	Percentage moisture content of packaged and stored <i>S. aethiopicum</i>					
	RC-CR			RC-CC		
	RC-CR0.1	RC-CR0.5	RC-CR60 μm	RC-CC0.1	RC-CC0.5	RC-CC60 μm
Zero	86.90 ± 1.76	86.91 ± 1.76	86.91 ± 1.76	86.91 ± 1.76	86.91 ± 1.76	86.91 ± 1.76
One	85.22 ± 0.06	86.03 ± 0.19	87.23 ± 0.19	86.55 ± 0.37	85.86 ± 0.43	85.37 ± 0.27
Two	84.88 ± 0.04	84.76 ± 0.09	86.17 ± 0.21	86.02 ± 0.19	85.39 ± 0.06	88.08 ± 0.24
Three	83.93 ± 0.19	84.62 ± 0.17	87.04 ± 0.51	86.62 ± 0.01	82.12 ± 0.47	86.48 ± 0.23
Four	82.99 ± 0.03	83.31 ± 0.10	85.66 ± 0.06	85.20 ± 0.05	80.55 ± 0.07	84.55 ± 0.18

Values are expressed as means \pm standard deviation. ^{abcde}Values not sharing common superscript within the same row are significantly different ($P \leq 0.05$) using Tukey HSD test. RC-CR is cold room and RC-CC is charcoal cooler. RC-CR0.1, RC-CR0.5 and RC-CR60 μm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μm perforated polyethylene respectively and kept in cold room. RC-CC0.1, RC-CC0.5 and RC-CC60 μm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μm perforated polyethylene respectively and kept in charcoal cooler.

The percentage weight loss of *S. aethiopicum* RC-CR0.1 and RC-CC0.1 increased from 0.00±0.00 % to 11.33±0.23% and 0.00±0.00 % to 12.57±0.27 % respectively. For the *S. aethiopicum* RC-CR0.5 and RC-CC0.5 the percentage weight loss increased from 0.00±0.00 % on day of harvest to 9.80±0.18 % and from 0.00±0.00 % to 12.49±0.05 % respectively as that of *S. aethiopicum* RC-CR60µm increased from 0.00±0.00 % to 8.83±0.72 % from 0.00±0.00 % to 9.69±0.25 % respectively. *S. aethiopicum* RC-CR0.1 and RC-CC0.1

showed the highest increase in percentage weight loss as RC-CR60µm perforated polyethylene showed the lowest percentage weight loss.

These results show that there was a faster increase in weight loss of packaged *S. aethiopicum* stored in charcoal cooler and also there was a significant difference (p≤0.05) in the percentage weight loss of the packaged *S. aethiopicum* stored in cold room and charcoal cooler for each day among the three different packaging materials.




Figure 9. Effect on chlorophyll content, of immersing harvested *S. aethiopicum* in portable water intermittently but stored at ambient condition (TT⁺) and that stored in charcoal cooler but without water treatment (TT⁻). RI; with roots intact and RC; with roots cut-off. RI(TT⁺)-AC and RC(TT⁺)-AC is *S. aethiopicum* in ambient storage with water treatment but with roots intact and roots cut-off respectively. RI(TT⁻)-CC and RC(TT⁻)-CC is *S. aethiopicum* in charcoal cooler without water treatment but with roots intact and roots cut-off respectively. Effect on chlorophyll content, of immersing harvested *S. aethiopicum* in portable water intermittently but stored at ambient condition (TT⁺) and that stored in charcoal cooler but without water treatment (TT⁻). RI; with roots intact and RC; with roots cut-off. RI(TT⁺)-AC and RC(TT⁺)-AC is *S. aethiopicum* in ambient storage with water treatment but with roots intact and roots cut-off respectively. RI(TT⁻)-CC and RC(TT⁻)-CC is *S. aethiopicum* in charcoal cooler without water treatment but with roots intact and roots cut-off respectively

Table 2. Percentage weight loss of packaged *S. aethiopicum* stored in cold room and charcoal cooler

Duration (days)	Percentage Weigh loss of packaged <i>S.aethiopicum</i> in different storage condition					
	RC-CR			RC-CC		
	RC-CR0.1	RC-CR0.5	RC-CR60µm	RC-CC0.1	RC-CC0.5	RC-CC60µm
Zero	0.00 ± 0.00	0.00 ± .00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
One	4.59 ± 0.25	4.44 ± 0.22	3.30 ± 0.81	5.21 ± 0.10	3.37 ± 1.24	3.27 ± 0.47
Two	5.24 ± 0.63	9.06 ± 0.10	5.04 ± 0.81	6.71 ± 0.21	4.03 ± 0.25	5.40 ± 0.12
Three	7.86 ± 0.28	9.34 ± 0.23	6.60 ± 1.07	10.07 ± 0.59	7.40 ± 0.16	7.73 ± 0.07
Four	11.33± 0.23	9.80 ± 0.18	8.83 ± 0.72	12.57 ± 0.27	12.49 ± 0.05	9.69 ± 0.25

Values are expressed as means ± standard deviation. Values within the same row and column are significantly different (P ≤ 0.05) using Tukey HSD test. RC-CR is cold room and RC-CC is charcoal cooler. RC-CR0.1, RC-CR0.5 and RC-CR60µm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 µm perforated polyethylene respectively and kept in cold room. RC-CC0.1, RC-CC0.5 and RC-CC60µm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 µm perforated polyethylene respectively and kept in charcoal cooler.

Table 3. Chlorophyll content of packaged *S.aethiopicum* stored in cold room and charcoal cooler

Duration (days)	Chlorophyll content (mg/g dwb) of packaged <i>S.aethiopicum</i> in different Storage conditions					
	RC-CR			RC-CC		
	RC-CR0.1	RC-CR0.5	RC-CR60µm	RC-CC0.1	RC-CC0.5	RC-CC60µm
Zero	19.88±0.03	19.88±0.03	19.88±0.03	19.88±0.03	19.88±0.03	19.88±0.03
One	18.30±0.36	20.87±0.01	18.80±0.06	19.24±0.07	16.69±0.14	19.52±0.01
Two	17.58±0.23	17.88±0.00	18.80±0.51	17.97±0.02	15.75±0.02	18.68±0.01
Three	17.07±0.01	15.36±0.16	20.12±0.14	14.42±0.09	8.02±0.06	13.57±0.04
Four	14.67±0.02	14.70±0.09	19.21±0.08	7.16±0.04	5.27±0.01	8.06±0.02

Values are expressed as means ± standard deviation. Values within the same row and column are significantly different (P ≤ 0.05) using Tukey HSD test. RC-CR is cold room and RC-CC is charcoal cooler. RC-CR0.1, RC-CR0.5 and RC-CR60µm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 µm perforated polyethylene respectively and kept in cold room. RC-CC0.1, RC-CC0.5 and RC-CC60µm is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 µm perforated polyethylene respectively and kept in charcoal cooler.

The chlorophyll content of *S. aethiopicum* RC-CR0.1 and RC-CR0.5 decreased from 19.88 ± 0.03 mg/g dwb to 14.67 ± 0.02 mg/g dwb and 14.70 ± 0.09 mg/g dwb respectively as shown in Table 3.

The chlorophyll content of *S. aethiopicum* decreased from 19.88 ± 0.03 mg/g dwb on day of harvest to 19.21 ± 0.08 for RC-CR60 μ m and to 8.06 ± 0.02 mg/g dwb for RC-CR60 μ m. *S. aethiopicum* RC-CC0.5 showed the most rapid decrease in chlorophyll content from 19.88 ± 0.03 mg/g dwb to 5.27 ± 0.01 mg/g dwb and *S. aethiopicum* RC-CR0.1 showed the highest decrease in chlorophyll content followed by *S. aethiopicum* RC-CR0.5 whereas RC-CR60 μ m lowest decrease in chlorophyll content. Packaged *S. aethiopicum* stored in charcoal cooler showed a more rapid decrease in chlorophyll compared to that stored in cold room for all the three packaging materials. There was a variation between the chlorophyll content of packaged *S. aethiopicum* in the three different packaging materials and within the storage condition which was significant at $p \leq 0.05$.

The polyphenol content of *S. aethiopicum* RC-CR0.1, RC-CR0.5 and RC-CR60 μ m increased from 3.30 ± 0.21 mgQE/gfw on day zero to 5.48 ± 0.03 mgQE/gfw, 5.52 ± 0.13 mgQE/gfw and 5.55 ± 0.09 mgQE/gfw on day four respectively. The polyphenol content of *S. aethiopicum* RC-CC0.1, RC-CC0.5 and RC-CC60 μ m increased from 3.30 ± 0.21 mgQE/gfw on day zero to 5.89 ± 0.01 mgQE/gfw, 4.13 ± 0.03 mgQE/gfw and 4.55 ± 0.05 mgQE/gfw on day four respectively.

The total polyphenol content increased with storage duration in both cold room and charcoal cooler for all the three packaging materials. The result analysis showed that there was a significant difference ($P \leq 0.05$) in the total polyphenol content with in the three different packaging

materials and storage conditions except on day four the RC-CR0.5 and RC-CR60 μ m also RC-CC0.5 and RC-CC60 μ m showed no significant difference ($P \leq 0.05$) as shown in Table 4

The TAA of *S. aethiopicum* RC-CR0., RC-CR0.5 and RC-CR60 μ m increased from 1.32 ± 0.18 mgAAE/gfw on day zero to 1.82 ± 0.03 mgAAE/gfw, 1.97 ± 0.15 mgAAE/gfw and 2.05 ± 0.02 mgAAE/gfw on day four respectively and RC-CC0., RC-CC0.5 and RC-CC60 μ m increased from 1.32 ± 0.18 mgAAE/gfw on day zero to 2.47 ± 0.02 mgAAE/gfw, 2.12 ± 0.15 mgAAE/gfw and 2.11 ± 0.02 mgAAE/gfw on day four respectively when using FRAP method. When using DPPH method, the TAA of *S. aethiopicum* packaged RC-CR0.1, RC-CR0.5 and RC-CR60 μ m increased from 2.02 ± 0.02 mgAAE/gfw on day zero to 2.73 ± 0.03 mgAAE/gfw, 2.78 ± 0.03 mgAAE/gfw and 3.13 ± 0.03 mgAAE/gfw on day four respectively while RC-CR0.1, RC-CR0.5 and RC-CR60 μ m increased from 2.02 ± 0.02 mgAAE/gfw on day zero to 3.73 ± 0.02 mgAAE/gfw, 3.20 ± 0.05 mgAAE/gfw and 2.89 ± 0.05 mgAAE/gfw on day four respectively. The TAA of the packaged *S. aethiopicum* increased with storage duration from day of harvest to day three and then slightly decreased, when determined using both FRAP and DPPH methods, for all the three different packaging materials for *S. aethiopicum* stored in cold room as shown in Figure 10.

The TAA of that stored in charcoal cooler generally increased with storage duration from day of harvest to day four when packaged in the three different packaging materials as shown in Figure 10. There was a variation between the TAA of packaged *S. aethiopicum* in the three different packaging materials and within the storage condition which was significant at $p \leq 0.05$.




Figure 10. Variation in the total antioxidant activity (TAA) of packaged *S. aethiopicum* and stored in cold room (CR) and charcoal cooler (CC). CR-DPPH and CC-DPPH is TAA of *S. aethiopicum* stored in CR and CC respectively when determined using DPPH method. CR-FRAP and CC-FRAP is TAA of *S. aethiopicum* stored in CR and CC respectively when determined using FRAP method. RC-CR0.1, RC-CR0.5 and RC-CR60 μ m is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μ m perforated polyethylene respectively and kept in cold room. RC-CC0.1, RC-CC0.5 and RC-CC60 μ m is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μ m perforated polyethylene respectively and kept in charcoal cooler

Table 4. Total polyphenol (T.P) content of packaged *S. aethiopicum* stored in cold room and charcoal cooler

Duration (days)	T.p content (mgQE/gfw) of packaged <i>S. aethiopicum</i> stored in cold room and charcoal cooler					
	RC-CR			RC-CC		
	RC-CR0.1	RC-CR0.5	RC-CR60 μ m	RC-CC0.1	RC-CC0.5	RC-CC60 μ m
Zero	3.30 \pm 0.21 ^a	3.30 \pm 0.21 ^a	3.30 \pm 0.21 ^a	3.30 \pm 0.21 ^a	3.30 \pm 0.21 ^a	3.30 \pm 0.21 ^a
One	3.20 \pm 0.05 ^a	3.73 \pm 0.03 ^b	3.43 \pm 0.03 ^c	3.92 \pm 0.05 ^d	4.52 \pm 0.08 ^e	3.63 \pm 0.03 ^f
Two	4.75 \pm 0.01 ^a	5.63 \pm 0.03 ^b	4.68 \pm 0.03 ^c	5.24 \pm 0.04 ^d	5.10 \pm 0.05 ^e	4.26 \pm 0.05 ^f
Three	6.10 \pm 0.06 ^a	4.90 \pm 0.05 ^b	5.08 \pm 0.14 ^c	5.61 \pm 0.01 ^d	4.88 \pm 0.03 ^e	5.61 \pm 0.05 ^f
Four	5.48 \pm 0.03 ^a	5.52 \pm 0.13 ^b	5.55 \pm 0.09 ^b	5.89 \pm 0.01 ^c	4.13 \pm 0.03 ^d	4.55 \pm 0.05 ^d

Values are expressed as means \pm standard deviation. ^{abcde}Values not sharing common superscript with in a row are significantly different ($P \leq 0.05$) using Tukey HSD test. RC-CR is cold room and RC-CC is charcoal cooler. RC-CR0.1, RC-CR0.5 and RC-CR60 μ m is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μ m perforated polyethylene respectively and kept in cold room. RC-CC0.1, RC-CC0.5 and RC-CC60 μ m is *S. aethiopicum* with roots cut-off and packaged in 0.1 cm meshed perforated polyethylene, 0.5 cm meshed perforated polyethylene, 60 μ m perforated polyethylene respectively and kept in charcoal cooler

4. Discussion

There are several factors that affect post-harvest quality of a vegetable, these include temperature in storage room, percentage relative humidity, light intensity, oxygen concentration, and carbon dioxide concentration [9]. The first indicator of deterioration of vegetables is moisture loss due to low relative humidity and high temperature, weight loss due to moisture loss and yellowing due to ethylene production [27]. The moisture loss is indicated by wilting, discolouration, and shriveling due to loss of turgidity and firmness [28]. The moisture content of the stored *S. aethiopicum* decreased with storage duration due to evaporation [29] resulting from fast air movement, low moisture content in the air and the thin waxy skin with numerous pores on the leaves. In the current study, the vegetables were subjected to different storage conditions with different temperatures and relative humidity and the findings indicated a difference in the moisture content in the vegetables stored under different conditions. Storage in the cold room and charcoal cooler showed higher retention of moisture compared to ambient storage which agrees with other studies [30,31]. This finding suggest that the lower the temperatures and the higher the relative humidity (experienced in both cold storage and charcoal cooler technology), the lower the rate of moisture loss and evaporation. Different vegetables respond to temperature variations after harvest and in storage differently [32]. Most of them however require a combination of low temperature ranging from 0°C to 10°C and high percentage relative humidity ranging from 85 % to 98 % [33]. This relative humidity can be achieved in either cold room or charcoal cooler. The low relative humidity below the optimum and high temperatures experienced in the ambient storage often lead to a higher rate of moisture loss. The rapid reduction in moisture content is seen to have caused a rapid increase in percentage weight loss (Figure 2). These agree with the findings reported elsewhere [34], confirming that any decrease in percentage moisture content leads in increase in percentage weight loss.

There was a decline in chlorophyll content with longer duration of storage of the leafy vegetable in all the three storage conditions. However, storage in ambient conditions revealed a faster decline most probably due to the high light intensity, leading to chlorophyll degradation and production of yellowish pigment as reported [35,36,37,38]. The light intensity in the charcoal cooler and cold room was relatively regulated because the vegetables were kept closed inside the structures and; particularly for the charcoal cooler, the inner walls were

lined with black cloth which naturally regulates light. Such a mechanism can reduce the rate of chlorophyll degradation. Additionally, the decline in chlorophyll content was more rapid for the *S. aethiopicum* with roots cut-off. It is postulated that it could be due to the increased production of ethylene levels resulting from wounding [39,40]. The high temperatures in ambient storage are known to cause an increase in the activity of polyphenol oxidase whose optimum temperature is 30°C [41] and its activity lead to browning.

Polyphenols are important chemicals synthesized by plants for protection especially from harsh conditions like water stress, high temperatures and wounding [10,11,42,43,44]. This study established that ambient storage has harsh conditions for storage for example high temperature and low relative humidity which are known causes of high polyphenol content in stored *S. aethiopicum*. The levels are not comparable to those under cold room and charcoal cooler storage conditions, irrespective of whether or not they have roots. The high light intensity in the ambient storage has also been reported to lead to increased polyphenol content in stored vegetables [35,45]. In this current study, the polyphenol content of *S. aethiopicum* in ambient decreased after day three, which could be attributed to the increase in the polyphenol oxidase activity [41]. In this study, the total polyphenol content of *S. aethiopicum* stored in cold room and charcoal cooler remained almost constant most likely due to the low temperature and high relative humidity. This reduced the moisture loss and resulted in regulation of enzyme activity. The increase in total polyphenol of *S. aethiopicum* with roots cut-off stored in ambient storage was faster, probably because of cutting which resulted in wounding of the vegetable leading to increase in total polyphenol content [46]. This was controlled in cold room and charcoal cooler storage conditions. The increase in the total antioxidant activity can be attributed to the increase in the total polyphenol content with storage duration. The positive linear correlation between total antioxidants and polyphenol content has been shown in several studies [47,48,49,50,51,52,53].

When the vegetable in ambient storage was intermittently immersed in portable water, a practice being carried out in markets to prolong shelf life, the total polyphenol and chlorophyll content remained almost constant probably due to cooling that reduced the enzyme activity for the biosynthesis of polyphenols and ethylene. This also increased the moisture on the leaf surface, maintaining a high relative humidity around the leaves leading to reduced evaporation [54]. Packaging prolongs

shelf life [55,56,57,58] and different packaging materials give different shelf life and quality of stored vegetables [56,57,59]. The 60 µm perforated polyethylene showed higher moisture retention. This can be attributed to the increased relative humidity in the inside [60] as the polyethylene does not allow water penetration [58]. The perforation reduces water vapour condensation [54] hence maintaining a high relative humidity inside the material and reducing water loss from the vegetable [54]. The polyethylene material has been reported to be good materials for decreasing physicochemical changes and nutrient quality changes [14,61]. Packaging is reported to reduce chlorophyll loss from vegetables [12]. The loss in chlorophyll content was more in the 0.1 cm meshed perforated polyethylene and 0.5 cm meshed perforated polyethylene. This is probably due to the large pore size which increase light transmission and oxygen concentration within the packaging material. This increased light transmission and oxygen concentration increases chlorophyll loss leading to deterioration [12,13,14].

The total polyphenol content of *S. aethiopicum* increased in all the packaging material. probably due to the post-harvest biochemical reactions that set in with moisture loss and wounding, resulting from the cutting off of roots and the roughness of the packaging material, that promotes phytochemical synthesis [10,11]. This increase in the total polyphenol content also lead to the increase in the total antioxidant activity of *S. aethiopicum* in all packaging materials as shown in Figure 10. The best packaging material studied for storage of *S. aethiopicum* was the 60 µm perforated polyethylene because of its ability to reduce moisture loss, reducing air and oxygen saturation inside the packaging hence probably increasing carbon dioxide concentration inside the packaging. This material is also smooth; this prevents wounding the leaves of the packaged vegetable. a proper packaging material should not be rough to avoid injuring the packaged vegetable contained inside as injuring a vegetable reduces the shelf life of the leafy vegetable [62].

5. Conclusion and Recommendations

Deterioration of harvested *S. aethiopicum* occurs more rapidly when the roots of the vegetable are cut-off. The percentage relative humidity is the most prominent factor that affected the moisture content and weight of stored *S. aethiopicum*. Cutting off of roots leads to increase in polyphenol biosynthesis and chlorophyll loss with storage duration as a result of increase in ethylene production of the vegetable leaves. The total antioxidant activity of *S. aethiopicum* increases with storage duration, however the quality of the vegetable in terms of moisture content, weight, nutritional quality has reduced by day four. Therefore, the consumer ought to consume this vegetable in the first four days if it is to be fed on when fresh. The freshness of the vegetable can be maintained by immersing the leaves intermittently in portable water for the first two to three days for vegetables in ambient storage however due to a lot of moisture, oxygen and microorganisms in the environment, rotting sets in and the quality of the vegetable reduces sharply. Therefore, the

stored *S. aethiopicum* must be kept away from contact with water to prevent rotting. This makes the charcoal cooler a better affordable storage technology as it allows no contact of the vegetable with water yet it maintains a high relative humidity required. Packaging of *S. aethiopicum* prolongs self life and 60 µm perforated polyethylene packaging material is a preferred for *S. aethiopicum*.

Acknowledgements

The authors are grateful for the research facilities offered by Makerere University School of Food Technology, Nutrition and Bio-engineering system, financial support provided by EU (PAEPARD/CRFII) through FARA and Uganda Christian University.

References

- [1] Chinedu, S.N., et al., Proximate and phytochemical analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. fruits. *Research Journal of Chemical Sciences*, 2011. 1(3): p. 63-71.
- [2] Ssekabembe, C., C. Bukenya, and W. Nakyagaba. Traditional knowledge and practices in local vegetable production in central Uganda. in *Afr Crop Sci Conf Proc*. 2003.
- [3] Naggayi, G., Vegetable Consumption and selected Micronutrient retention of Prepared and Cooked *Solanum aethiopicum* (Shum). A case for Rubaga division, Kampala., in A dissertation submitted to the College of Agricultural and Environmental sciences. 2016, Makerere University.
- [4] El-Ramady, H.R., et al., Postharvest management of fruits and vegetables storage, in *Sustainable agriculture reviews*. 2015, Springer. p. 65-152.
- [5] Prabhu, S. and D.M. Barrett, Effects of storage condition and domestic cooking on the quality and nutrient content of African leafy vegetables (*Cassia tora* and *Corchorus tridens*). *Journal of the Science of Food and Agriculture*, 2009. 89(10): p. 1709-1721.
- [6] Horna, D., S. Timpo, and G. Gruère, Marketing underutilized crops: The case of the African garden egg (*Solanum aethiopicum*) in Ghana. *Global Facilitation Unit for Underutilized Species (GFU) Via dei Tre Denri, Rome, Italy, 2007*: p. 3.
- [7] Nakanwagi, M.J., et al., Performance of *Solanum aethiopicum* Shum group accessions under repetitive drought stress. *Journal of Plant Breeding and Crop Science*, 2018. 10(1): p. 13-20.
- [8] Apolot, M.G., Profile and estimates of postharvest losses that occur along value chains of *Solanum aethiopicum* shum (nakati) and *Amaranthus lividus* (bugga) leafy vegetables, in A dissertation submitted to the college of Agriculture and Environmental Sciences. 2018, Makerere University: Makerere University 2018.
- [9] Shivashankara, K.S., N.K.S. Rao, and G.A. Geetha, Impact of climate change on fruit and vegetable quality, in *Climate-resilient Horticulture: Adaptation and Mitigation Strategies*. 2013, Springer. p. 237-244.
- [10] Khlestkina, E., The adaptive role of flavonoids: emphasis on cereals. *Cereal Research Communications*, 2013. 41(2): p. 185-198.
- [11] Park, S., et al., Determination of polyphenol levels variation in *Capsicum annum* L. cv. Chelsea (yellow bell pepper) infected by anthracnose (*Colletotrichum gloeosporioides*) using liquid chromatography–tandem mass spectrometry. *Food Chemistry*, 2012. 130(4): p. 981-985.
- [12] Pristouri, G., A. Badeka, and M. Kontominas, Effect of packaging material headspace, oxygen and light transmission, temperature and storage time on quality characteristics of extra virgin olive oil. *Food control*, 2010. 21(4): p. 412-418.
- [13] Sahoo, N.R., et al., A comparative study on the effect of packaging material and storage environment on shelf life of fresh bell-pepper. *Journal of Food Measurement and Characterization*, 2014. 8(3): p. 164-170.

- [14] Sahoo, N.R., et al., Effect of packaging conditions on quality and shelf-life of fresh pointed gourd (*Trichosanthes dioica* Roxb.) during storage. *Food Packaging and Shelf Life*, 2015. 5: p. 56-62.
- [15] Akanyijuka, S., Effect of different processing conditions on proximate and bioactive contents of *Solanum aethiopicum* (Shum) powders and Acceptability for cottage scale production., in A dissertation submitted to the college of Agricultural and Environmental Sciences. 2017, Makerere University.
- [16] Boyhan, G.E., et al., Sowing date, transplanting date, and variety effect on transplanted short-day onion production. *HortTechnology*, 2009. 19(1): p. 66-71.
- [17] Rickman, J.C., C.M. Bruhn, and D.M. Barrett, Nutritional comparison of fresh, frozen, and canned fruits and vegetables II. Vitamin A and carotenoids, vitamin E, minerals and fiber. *Journal of the Science of Food and Agriculture*, 2007. 87(7): p. 1185-1196.
- [18] Wissam, Z., et al., Effective extraction of polyphenols and proanthocyanidins from pomegranate's peel. *Int J Pharm Pharm Sci*, 2012. 4(Supplement 3): p. 675-682.
- [19] Demiray, S., M. Pintado, and P. Castro, Evaluation of phenolic profiles and antioxidant activities of Turkish medicinal plants: *Tilia argentea*, *Crataegi folium* leaves and *Polygonum bistorta* roots. *World Academy of Science, Engineering and Technology*, 2009. 54: p. 312-317.
- [20] Majhenič, L., M. Škerget, and Ž. Knez, Antioxidant and antimicrobial activity of guarana seed extracts. *Food chemistry*, 2007. 104(3): p. 1258-1268.
- [21] Braca, A., et al., Antioxidant principles from *bauhinia tarapotensis*. *Journal of natural products*, 2001. 64(7): p. 892-895.
- [22] Iqbal, E., K.A. Salim, and L.B. Lim, Phytochemical screening, total phenolics and antioxidant activities of bark and leaf extracts of *Goniothalamus velutinus* (Airy Shaw) from Brunei Darussalam. *Journal of King Saud University-Science*, 2015. 27(3): p. 224-232.
- [23] Lu, Y., T. Khoo, and C. Wiart, Antioxidant activity determination of citronellal and crude extracts of *Cymbopogon citratus* by 3 different methods. *Pharmacology & Pharmacy*, 2014. 5(04): p. 395.
- [24] Sumanta, N., et al., Spectrophotometric analysis of chlorophylls and carotenoids from commonly grown fern species by using various extracting solvents. *Research Journal of Chemical Sciences ISSN*, 2014. 2231: p. 606X.
- [25] Lichtenthaler, H.K. and C. Buschmann, Chlorophylls and carotenoids: Measurement and characterization by UV-VIS spectroscopy. *Current protocols in food analytical chemistry*, 2001.
- [26] AOAC, Official Methods of Analysis of the Association of Official Analytical Chemists. 1995, AOAC: Arlington, Virginia.
- [27] Karaman, S., et al., A response surface methodology study on the effects of some phenolics and storage period length on vegetable oil quality: change in oxidation stability parameters. *Turkish Journal of Agriculture and Forestry*, 2014. 38(6): p. 759-772.
- [28] Ambuko, J., et al., Preservation of Postharvest Quality of Leafy Amaranth (*Amaranthus* spp.) Vegetables Using Evaporative Cooling. *Journal of Food Quality*, 2017. 2017.
- [29] Snyder, K., J. Richards, and L. Donovan, Night-time conductance in C3 and C4 species: do plants lose water at night? *Journal of Experimental Botany*, 2003. 54(383): p. 861-865.
- [30] Krokida, M.K., et al., Drying kinetics of some vegetables. *Journal of Food engineering*, 2003. 59(4): p. 391-403.
- [31] Gawrysiak-Witulska, M., et al., The effect of temperature and moisture content of stored rapeseed on the phytosterol degradation rate. *Journal of the American Oil Chemists' Society*, 2012. 89(9): p. 1673-1679.
- [32] do Nascimento Nunes, M.C., Impact of environmental conditions on fruit and vegetable quality. *Stewart Postharvest Review*, 2008. 4(4).
- [33] Thompson, J., A. Kader, and K. Sylva, Compatibility chart for fruits and vegetables in short-term transport or storage. *Oakland: Univ. Calif. Div. Ag. and Nat. Res. Publ.* 1996. 21560.
- [34] Kwenin, W., M. Wollin, and B. Dzomeku, Assessing the nutritional value of some African indigenous green leafy vegetables in Ghana. *Journal of Animal and Plant Sciences*, 2011. 10(2): p. 1300-1305.
- [35] Jin, P., et al., Effect of light on quality and bioactive compounds in postharvest broccoli florets. *Food chemistry*, 2015. 172: p. 705-709.
- [36] He, Q., et al., Effect of light intensity on physiological changes, carbon allocation and neutral lipid accumulation in oleaginous microalgae. *Bioresource technology*, 2015. 191: p. 219-228.
- [37] Bielczynski, L.W., G. Schansker, and R. Croce, Effect of light acclimation on the organization of photosystem II super- and sub-complexes in *Arabidopsis thaliana*. *Frontiers in plant science*, 2016. 7.
- [38] Gomes, M.P., et al., Differential effects of glyphosate and aminomethylphosphonic acid (AMPA) on photosynthesis and chlorophyll metabolism in willow plants. *Pesticide biochemistry and physiology*, 2016. 130: p. 65-70.
- [39] Brecht, J.K., Physiology of lightly processed fruits and vegetables. *HortScience*, 1995. 30(1): p. 18-22.
- [40] Saltveit, M. The three responses of plant tissue to wounding. in III International Conference on Fresh-Cut Produce: Maintaining Quality and Safety 1141. 2015.
- [41] Bello, A. and M. Sule, Optimum temperature and thermal stability of crude polyphenol oxidase from some common fruits. *Nigerian Journal of Basic and Applied Sciences*, 2012. 20(1): p. 27-31.
- [42] Hou, J., et al., Genome-wide transcriptomic profiles reveal multiple regulatory responses of poplar to *Lonsdalea quercina*. *Trees*, 2016. 30(4): p. 1389-1402.
- [43] Kim, H.G., et al., Determination of the change of flavonoid components as the defence materials of *Citrus unshiu* Marc. fruit peel against *Penicillium digitatum* by liquid chromatography coupled with tandem mass spectrometry. *Food Chemistry*, 2011. 128(1): p. 49-54.
- [44] Rusjan, D., et al., Biochemical response of grapevine variety 'Chardonnay' (*Vitis vinifera* L.) to infection with grapevine yellows (Bois noir). *European journal of plant pathology*, 2012. 134(2): p. 231-237.
- [45] Lee, E., H. Ahn, and E. Choe, Effects of light and lipids on chlorophyll degradation. *Food Science and Biotechnology*, 2014. 23(4): p. 1061-1065.
- [46] Toivonen, P.M. and D.A. Brummell, Biochemical bases of appearance and texture changes in fresh-cut fruit and vegetables. *Postharvest Biology and Technology*, 2008. 48(1): p. 1-14.
- [47] Stojadinovic, M., et al., Binding affinity between dietary polyphenols and β -lactoglobulin negatively correlates with the protein susceptibility to digestion and total antioxidant activity of complexes formed. *Food chemistry*, 2013. 136(3): p. 1263-1271.
- [48] Do, Q.D., et al., Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of food and drug analysis*, 2014. 22(3): p. 296-302.
- [49] Genskowsky, E., et al., Determination of polyphenolic profile, antioxidant activity and antibacterial properties of maqui [*Aristotelia chilensis* (Molina) Stuntz] a Chilean blackberry. *Journal of the Science of Food and Agriculture*, 2016. 96(12): p. 4235-4242.
- [50] Li, W., et al., Evaluation of Antioxidant Activity in Vitro of Polyphenols from *Toona sinensis* Seeds Using RSM Optimized Conditions. 2017.
- [51] Padhi, E.M., et al., Total polyphenol content, carotenoid, tocopherol and fatty acid composition of commonly consumed canadian pulses and their contribution to antioxidant activity. *Journal of Functional Foods*, 2017. 38: p. 602-611.
- [52] Podio, N.S., M.V. Baroni, and D.A. Wunderlin, Relation between polyphenol profile and antioxidant capacity of different Argentinean wheat varieties. A Boosted Regression Trees study. *Food Chemistry*, 2017. 232: p. 79-88.
- [53] Khan, I., et al., Acute effect of sorghum flour-containing pasta on plasma total polyphenols, antioxidant capacity and oxidative stress markers in healthy subjects: A randomised controlled trial. *Clinical Nutrition*, 2015. 34(3): p. 415-421.
- [54] Caleb, O.J., et al., Integrated modified atmosphere and humidity package design for minimally processed Broccoli (*Brassica oleracea* L. var. *italica*). *Postharvest Biology and Technology*, 2016. 121: p. 87-100.
- [55] Esturk, O., Z. Ayhan, and T. Gokkurt, Production and application of active packaging film with ethylene adsorber to increase the shelf life of broccoli (*Brassica oleracea* L. var. *italica*). *Packaging Technology and Science*, 2014. 27(3): p. 179-191.
- [56] Hailu, M., T.S. Workneh, and D. Belew, Effect of packaging materials on shelf life and quality of banana cultivars (*Musa* spp.). *Journal of food science and technology*, 2014. 51(11): p. 2947-2963.
- [57] Jacobsson, A., T. Nielsen, and I. Sjöholm, Effects of type of packaging material on shelf-life of fresh broccoli by means of changes in weight, colour and texture. *European Food Research and Technology*, 2004. 218(2): p. 157-163.

- [58] Udomkun, P., et al., Compositional and functional dynamics of dried papaya as affected by storage time and packaging material. *Food chemistry*, 2016. 196: p. 712-719.
- [59] Jaggi, M.P., R. Sharma, and S. Vali, Effect of packaging and storage conditions on retention of ascorbic acid in fenugreek leaves (*Trigonella Foenum-Graecum*). *Ukrainian Food Journal*, 2015. 4(2).
- [60] Schreiner, M., et al., Effect of film packaging and surface coating on primary and secondary plant compounds in fruit and vegetable products. *Journal of food engineering*, 2003. 56(2): p. 237-240.
- [61] Hymavathi, T. and V. Khader, Carotene, ascorbic acid and sugar content of vacuum dehydrated ripe mango powders stored in flexible packaging material. *Journal of Food Composition and Analysis*, 2005. 18(2): p. 181-192.
- [62] Ariffin, S.H., K. Gkatzionis, and S. Bakalis, Leaf injury and its effect towards shelf-life and quality of ready-to-eat (RTE) spinach. *Energy Procedia*, 2017. 123: p. 105-112.